

RECOMMENDED IMMUNISATION SCHEDULE

Vaccines	Protects Against	Recommended Age	Side Effects
BCG	Tuberculosis	At birth or within 6 weeks	Nodule formation after 3-6 wks. Ulcer with pus discharge. Scar by 8-12 wks.
DaPT, Hep B, Hib, IPV (Hexavalent) and OPV I	Diphtheria, Pertusis, Tetanus, Hepatitis B, Hemophilus Influenzae Meningitis & Pneumonia, Polio	1 st dose: 6 weeks (One and half months)	Slight pain & fever
DaPT, Hep B, Hib, IPV (Hexavalent) and OPV II		2 nd dose: 10 weeks (Two and half months)	
DaPT, Hep B, Hib, IPV (Hexavalent) and OPV III		3 rd dose: 14 weeks (Three and half months)	
Rotavirus I	Rotaviral Diarrhea	6 Weeks	No reaction. Rarely intussusception.
Rotavirus II		10 Weeks	
Rotavirus III		14 Weeks	
Pneumococcal 13 I	Pneumococcal Meningitis & Pneumonia	6 Weeks	Slight pain & fever
Pneumococcal 13 II		10 Weeks	
Pneumococcal 13 III		14 Weeks	
Pneumococcal 13 Booster		12 - 18 months	
Influenza Vaccine	Flu	6 months, 7 months and then yearly	Slight pain & fever
Typhoid	Typhoid	6 months & every 3 years	Mild fever
Measles-Mumps-Rubella (M.M.R.)	Measles, Mumps & Rubella	9 - 12 months	Mild rash & fever may appear after one week
Meningococcal ACWY	Meningococcal Meningitis	9 months & 2 nd dose three months after the 1 st dose	Slight pain & fever
Japanese Encephalitis	Japanese Encephalitis	12 - 23 months	Slight pain & fever
Hepatitis A I	Hepatitis A	1 year (Only Single dose for live vaccine)	Slight pain & fever
Hepatitis A II		18 - 24 months	
Chickenpox	Varicella	1 year and every 3 years	Mild rash & fever
Measles-Mumps-Rubella		15 - 18 months	
DPT, Hep B, HiB Booster		18 - 24 months	
M.M.R. Booster		4 - 5 years	
DaPT Booster		4 - 5 years	
M.M.R. Booster		9 - 10 years	
DaPT Booster		9 - 10 years	
Human Papilloma Virus HPV (For Girls)	Cervical, Vulvar & Vaginal Cancer	9 - 14 years (Two doses at gap of 6 months) 14 - 45 years (Three doses at 0, 2, and 6 months)	Slight pain & fever

All of the above mentioned vaccines should be given after one to one discussion with the parents.